

1995 Abstract Form  
Scientific Presentations  
F MAGNETIC RESONANCE  
SCIENTIFIC MEETING

N SOCIETY FOR MAGNETIC  
IN MEDICINE AND BIOLOGY  
TH ANNUAL MEETING

ropolis, NICE, FRANCE  
August 19-25, 1995

Presentation but  
sent as a poster  
but willing to make  
ation

quired (available only  
presentations)

GORIES  
OULD FILL IN A  
CATEGORY

/RY ☐  
ual Studies  
man White Matter  
man Other  
Neuro—Clinical  
Spine and Other  
lar  
amics and Flow  
lar—Clinical  
t  
inal  
ry  
age, Bone and Marrow  
aging  
al MRI

OPY  
RY ☐  
ic Imaging Techniques  
Spectroscopy  
Quantitation  
in—White Matter & Neuro  
e  
n—Stroke & Seizure  
n—Other  
n  
lar

letal  
ethods and Animal Models  
inical  
ding body fluids)

OGY

RY ☒

ification

Neuro—Methodology and Analysis  
Neuro—Models and Mechanisms  
Non-proton Imaging and ESR  
d Hardware

ig  
ization/Imaging  
Artifacts  
nces/Techniques  
Imaging  
sing  
Rendering  
chanisms/MTC  
e Contrast Agents  
st Agents  
lects  
al MRI

DEADLINE:

r than April 11, 1995.  
accepted abstracts become the  
MR. No proprietary information  
by authors.

Magnetic Resonance  
entile Meeting  
1a Street, Suite 201  
CA 94704, USA

g abstracts from outside the  
ow six weeks for mailing or  
by express.

ame and complete  
f the first author.

A. Bandettini

1 - NMR Center

Building 149

MA 02129

A

17) 726-8790

7) 726-7422

☒ Yes ☐ No

Only:

## FMRI and PET Demonstrate Sustained Blood Oxygenation and Flow Enhancement During Extended Visual Stimulation Durations

P. A. Bandettini<sup>1</sup>, T. L. Davis<sup>1,2</sup>, K. K. Kwong<sup>1</sup>, P. T. Fox<sup>3</sup>, A. Jiang<sup>1</sup>, J. R. Baker<sup>1,2</sup>,  
J. W. Belliveau<sup>1</sup>, R. M. Weisskoff<sup>1</sup>, and B. R. Rosen<sup>1</sup>

<sup>1</sup>Massachusetts General Hospital - NMR Center, Department of Radiology, Charlestown, MA

<sup>2</sup>Harvard - MIT Division of Health Sciences and Technology, Cambridge, MA

<sup>3</sup>University of Texas Health Science Center, San Antonio, TX

### INTRODUCTION:

Characterization of activation - induced changes in blood flow and oxygenation across time scales is one avenue by which neuronal, metabolic, and hemodynamic relationships may be more fully understood. Prior studies (1) have suggested that local blood oxygenation returns from an initially elevated level to baseline level after about 15 minutes of continuous stimulation. Possible reasons for this observation include either a) decreased neuronal firing with habituation or fatigue or b) an increase in oxidative metabolic rate over time. If the first reason applies, both oxygenation and flow should return to resting levels. If the second reason applies, then oxygenation should decrease but flow should remain elevated.

To test these hypotheses, both blood oxygenation - sensitive (gradient-echo, T2\*-weighted) and flow - sensitive (2) (inversion recovery, T1-weighted) images were collected during extended duration visual stimulation. Also, using flow measurements obtained by positron emission tomography (PET), habituation effects in primary visual and somatosensory cortex were assessed.

### METHODS:

**MRI:** Five subjects were studied. Flow sensitive (inversion recovery EPI: TI = 1000ms, TR = 3000ms, TE=20ms: spin-echo) and blood oxygenation sensitive (gradient-echo EPI: TR = 3000ms, TE = 40ms) time course series were obtained using a 1.5 T GE Signa scanner retrofitted with an ANMR resonant gradient system. Voxel volume was 3.125 mm x 3.125 mm x 15 mm. Motion correction (3), performed to reduce signal drift related to subject movement, improved signal stability in several experimental runs. Visual stimulation was: 10 Hz full field black and white alternating checkerboard. The timing was: 1 min.off, 1 min.on, 1 min.off, 20 min.on, 1 min.off, 1 min.on, 1 min.off.

**PET:** Seven and nine subjects were used in the visual stimulation and vibrotactile studies respectively. Cerebral blood flow was measured 40 seconds after bolus intravenous injections of <sup>15</sup>O-labeled water at 10 minute intervals during 50 minutes of either continuous 8 Hz full field alternating red and black checkerboard stimulation or vibrotactile stimulation.

### RESULTS:

Figures 1 and 2 show averaged, from activated visual cortex of 5 subjects, oxygenation-sensitive and flow-sensitive time course series respectively. No subject demonstrated a significant decrease in flow or oxygenation during the 20 minute stimulation period. A post-stimulation undershoot was also observed with the T2\*-weighted sequence and not the T1-weighted sequence.

Figure 3 shows averaged PET measurements, across all subjects. With visual stimulation, the activation-induced flow change did not decrease significantly. With vibrotactile stimulation, the flow change did decrease significantly over time. These results demonstrate that the degree of habituation may vary depending on the stimulus and activated region.

### CONCLUSIONS:

Blood flow and oxygenation in V1 remain elevated during extended stimulation. These studies agree with preliminary studies suggesting stable levels of oxidative metabolic rate, and oxygen extraction fraction in V1 during extended stimulation times (4,5). Reasons for the differences between the presented results and those of (1) may include differences in pulse sequence parameters

and/or differences in the type of visual stimulation used. Our visual stimulus may be: a) causing less habituation or neuronal fatigue and/or b) selectively activating neurons that do not increase as much in oxidative metabolic rate. Ongoing studies are being performed clarify these issues.




Figure 1: Averaged oxygenation - sensitive MR signal. GE-EPI: TR=3000ms. TE=40ms.




Figure 2: Averaged flow - sensitive MR signal. IR-EPI: TI=1000ms, TR=3000ms, TE=20ms: spin-echo.




Figure 3: Flow change in V1 and S1 over 50 minutes continual stimulation obtained using <sup>15</sup>O - labeled water injections at 10 minute intervals. A significant flow decrease is observed in S1 but not in V1, demonstrating that extended duration flow responses can vary depending on the stimuli used and cortical regions activated.

### REFERENCES:

1. Hathout GM, Kirlow KAT, So GJK, et al. *JMRI*, 4, 537-543 (1994).
2. Kwong KK, Belliveau JW, Chesler DA, et al. *PNAS*, 89, 5672, (1992).
3. Woods RP, Cherry SR, Mazziotta JC, *J. Comput. Assist Tomogr* 16, 620-633 (1992).
4. Fox PT, Personal communication.
5. Marrett S, Personal communication.